PAIN

Pain 101 (2003) 309–310

www.elsevier.nl/locate/pain

Author Index to Volume 101 (2003)

Aalto, S., see Hagelberg, N., **101** 149
Aglioti, S.M., see Rosso, T., **101** 117
Alexander, G.E., see Fukunaga, A.F., **101** 129
Arendt-Nielsen, L., see Svensson, P., **101** 221
Aziz, T., see Nandi, D., **101** 97

Bach, F.W., see Otto, M., **101** 187
Bak, S., see Otto, M., **101** 187
Barlas, P., see Chesterton, L.S., **101** 259
Basbaum, A.I., Editorial, **101** 1
Basbaum, A.I., see Malmberg, A.B., **101** 109
Baxter, G.D., see Chesterton, L.S., **101** 259
Beaudet, A., see Cahill, C.M., **101** 199
Beitz, A.J., see Wacnik, P.W., **101** 175
Bohotin, C., Scholsem, M., Multon, S., Martin, D., Bohotin, V., Schoenen, J., Vagus nerve stimulation in awake rats reduces formalin-induced nociceptive behaviour and fos-immunoreactivity in trigeminal nucleus caudalis, **101** 3
Bohotin, V., see Bohotin, C., **101** 3
Borchgrevink, P.C., see Klepstad, P., **101** 193
Bussmann, J.B.J., see Schasfoort, F.C., **101** 79

Cahill, C.M., Morinville, A., Hoffert, C., O'Donnell, D., Beaudet, A., Up-regulation and trafficking of δ opioid receptor in a model of chronic inflammation: implications for pain control, **101** 199
Cairns, B.E., see Svensson, P., **101** 221
Carpenter, R., see Eisenach, J.C., **101** 89
Carr, D.B., Cepeda, M.S., Reply to Letter to the Editor, **101** 211
Carter, H., see Nandi, D., **101** 97
Carter, P., see Hefferan, M.P., **101** 139
Cassidy, E.M., Tomkins, E., Sharifi, N., Dinan, T., Hardiman, O., O'Keane, V., Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
Cepeda, M.S., see Carr, D.B., **101** 211
Chang, V.T., see Hwang, S.S., **101** 55
Chesterton, L.S., Barlas, P., Foster, N.E., Baxter, G.D., Wright, C.C., Gender differences in pressure pain threshold in healthy humans, **101** 259
Curry, R., see Eisenach, J.C., **101** 89

de Tommaso, M., Valeriani, M., Guido, M., Libro, G., Specchio, L.M., Tonali, P., Puca, F., Abnormal brain processing of cutaneous pain in patients with chronic migraine, **101** 25
De, A., see Kimble, L.P., **101** 45
Devor, M., On dialog between clinicians and basic scientists, **101** 307
Dinan, T., see Cassidy, E.M., **101** 283
Dunbar, S.B., see Kimble, L.P., **101** 45
Dworkin, R.H., see Nagasako, E.M., **101** 213

Edwards, R.R., Fillingim, R.B., Ness, T.J., Age-related differences in endogenous pain modulation: a comparison of diffuse noxious inhibitory controls in healthy older and younger adults, **101** 155
Eisenach, J.C., Carpenter, R., Curry, R., Analgesia from a peripherally active κ-opioid receptor agonist in patients with chronic pancreatitis, **101** 89
Ernberg, M., Lundeberg, T., Kopp, S., Effects on muscle pain by intramuscular injection of granisetron in patients with fibromyalgia, **101** 275

doi:10.1016/S0304-3959(03)00034-4

Farina, S., see Rosso, T., **101** 117
Fazio, S., see Kimble, L.P., **101** 45
Fiaschi, A., see Rosso, T., **101** 117
Fillingim, R.B., see Edwards, R.R., **101** 155
Finco, G., see Rosso, T., **101** 117
Forssell, H., see Hagelberg, N., **101** 149
Foster, N.E., see Chesterton, L.S., **101** 259
Fukunaga, A.F., Alexander, G.E., Stark, C.W., Characterization of the analgesic actions of adenosine: comparison of adenosine and remifentanil infusions in patients undergoing major surgical procedures, **101** 129
Fukuoka, T., see Obata, K., **101** 65

Gilbert, H., see Malmberg, A.B., **101** 109
Graven-Nielsen, T., see Svensson, P., **101** 221
Gross, R.T., see Vowles, K.E., **101** 291
Gu, J.G., see Takeda, D., **101** 13
Guido, M., see de Tommaso, M., **101** 25

Häbler, H.-J., Jänig, W., Reflexes in sympathetic vasoconstrictor neurones arising from urinary bladder afferents are not amplified early after inflammation in the anaesthetised cat, **101** 251
Hagelberg, N., Forssell, H., Rinne, J.O., Scheinin, H., Taiminen, T., Aalto, S., Luutonen, S., Någren, K., Jääskeläinen, S., Striatal dopamine D1 and D2 receptors in burning mouth syndrome, **101** 149
Haley, M., see Hefferan, M.P., **101** 139
Hardiman, O., see Cassidy, E.M., **101** 283
Hashimoto, N., see Obata, K., **101** 65
Hazebroek-Kampschreur, A.A.J.M., see Merlijn, V.P.B.M., **101** 33
Hefferan, M.P., Carter, P., Haley, M., Loomis, C.W., Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia, **101** 139
Hoffert, C., see Cahill, C.M., **101** 199
Hollins, M., see Roy, E.A., **101** 267
Horowitz, S.H., Response to Letter, **101** 307
Hu, J.W., see Svensson, P., **101** 221
Hunfeld, J.A.M., see Merlijn, V.P.B.M., **101** 33
Hval, B., see Klepstad, P., **101** 193
Hwang, S.S., Chang, V.T., Kasimis, B., Cancer breakthrough pain characteristics and responses to treatment at a VA medical center, **101** 55

Ischia, S., see Rosso, T., **101** 117

Jääskeläinen, S., see Hagelberg, N., **101** 149
Jänig, W., see Häbler, H.-J., **101** 251
Jensen, T.S., see Otto, M., **101** 187
Jørum, E., Warncke, T., Stubhaug, A., Cold allodynia and hyperalgesia in neuropathic pain: the effect of *N*-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo, **101** 229
Jystad, Å., see Klepstad, P., **101** 193

Kaasa, S., see Klepstad, P., **101** 193
Kasimis, B., see Hwang, S.S., **101** 55
Kehl, L.J., see Wacnik, P.W., **101** 175

Kimble, L.P., McGuire, D.B., Dunbar, S.B., Fazio, S., De, A., Weintraub, W.S., Strickland, O.S., Gender differences in pain characteristics of chronic stable angina and perceived physical limitation in patients with coronary artery disease, **101** 45
Klepstad, P., Kaasa, S., Jystad, Å., Hval, B., Borchgrevink, P.C., Immediate- or sustained-release morphine for dose finding during start of morphine to cancer patients: a randomized, double-blind trial, **101** 193
Koes, B.W., see Merlijn, V.P.B.M., **101** 33
Kopp, S., see Ernberg, M., **101** 275

Libro, G., see de Tommaso, M., **101** 25
Loomis, C.W., see Hefferan, M.P., **101** 139
Lundeberg, T., see Ernberg, M., **101** 275
Luutonen, S., see Hagelberg, N., **101** 149

Maixner, W., see Roy, E.A., **101** 267
Malmberg, A.B., Gilbert, H., McCabe, R.T., Basbaum, A.I., Powerful antinociceptive effects of the cone snail venom-derived subtype-selective NMDA receptor antagonists conantokins G and T, **101** 109
Martin, D., see Bohotin, C., **101** 3
McCabe, R.T., see Malmberg, A.B., **101** 109
McCracken, L., see Roelofs, J., **101** 299
McGuire, D.B., see Kimble, L.P., **101** 45
Mehlisch, D.R., The combination of low dose of naloxone and morphine in patient-controlled (PCA) does not decrease opioid requirements in the postoperative period, **101** 209
Merlijn, V.P.B.M., Hunfeld, J.A.M., van der Wouden, J.C., Hazebroek-Kampschreur, A.A.J.M., Koes, B.W., Passchier, J., Psychosocial factors associated with chronic pain in adolescents, **101** 33
Morinville, A., see Cahill, C.M., **101** 199
Multon, S., see Bohotin, C., **101** 3

Nagasako, E.M., Oaklander, A.L., Dworkin, R.H., Congenital insensitivity to pain: an update, **101** 213
Någren, K., see Hagelberg, N., **101** 149
Nakatsuka, T., see Takeda, D., **101** 13
Nandi, D., Aziz, T., Carter, H., Stein, J., Thalamic field potentials in chronic central pain treated by periventricular gray stimulation – a series of eight cases, **101** 97
Ness, T.J., see Edwards, R.R., **101** 155
Noguchi, K., see Obata, K., **101** 65

Oaklander, A.L., see Nagasako, E.M., **101** 213
Obata, K., Yamanaka, H., Fukuoka, T., Yi, D., Tokunaga, A., Hashimoto, N., Yoshikawa, H., Noguchi, K., Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats, **101** 65
O'Donnell, D., see Cahill, C.M., **101** 199
O'Keane, V., see Cassidy, E.M., **101** 283
Otto, M., Bak, S., Bach, F.W., Jensen, T.S., Sindrup, S.H., Pain phenomena and possible mechanisms in patients with painful polyneuropathy, **101** 187

Papke, R., see Takeda, D., **101** 13
Passchier, J., see Merlijn, V.P.B.M., **101** 33
Pertovaara, A., Wei, H., A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Peters, M.L., see Roelofs, J., **101** 299
Price, D.D., see Staud, R., **101** 167
Puca, F., see de Tommaso, M., **101** 25

Ramnaraine, M.L., see Wacnik, P.W., **101** 175
Rinne, J.O., see Hagelberg, N., **101** 149
Robinson, M.E., see Staud, R., **101** 167
Roelofs, J., Peters, M.L., McCracken, L., Vlaeyen, J.W.S., The pain vigilance and awareness questionnaire (PVAQ): further psychometric evaluation in fibromyalgia and other chronic pain syndromes, **101** 299
Rosso, T., Aglioti, S.M., Zanette, G., Ischia, S., Finco, G., Farina, S., Fiaschi, A., Tinazzi, M., Functional plasticity in the human primary somatosensory cortex following acute lesion of the anterior lateral spinal cord: neurophysiological evidence of short-term cross-modal plasticity, **101** 117
Roy, E.A., Hollins, M., Maixner, W., Reduction of TMD pain by high-frequency vibration: a spatial and temporal analysis, **101** 267

Schasfoort, F.C., Bussmann, J.B.J., Zandbergen, A.M.A.J., Stam, H.J., Impact of upper limb complex regional pain syndrome type 1 on everyday life measured with a novel upper limb-activity monitor, **101** 79
Scheinin, H., see Hagelberg, N., **101** 149
Schoenen, J., see Bohotin, C., **101** 3
Scholsem, M., see Bohotin, C., **101** 3
Sessle, B.J., see Svensson, P., **101** 221
Sharifi, N., see Cassidy, E.M., **101** 283
Sindrup, S.H., see Otto, M., **101** 187
Specchio, L.M., see de Tommaso, M., **101** 25
Stam, H.J., see Schasfoort, F.C., **101** 79
Stark, C.W., see Fukunaga, A.F., **101** 129
Staud, R., Robinson, M.E., Vierck Jr, C.J., Price, D.D., Diffuse noxious inhibitory controls (DNIC) attenuate temporal summation of second pain in normal males but not in normal females or fibromyalgia patients, **101** 167
Stein, J., see Nandi, D., **101** 97
Strickland, O.S., see Kimble, L.P., **101** 45
Stubhaug, A., see Jørum, E., **101** 229
Svensson, P., Cairns, B.E., Wang, K., Hu, J.W., Graven-Nielsen, T., Arendt-Nielsen, L., Sessle, B.J., Glutamate-evoked pain and mechanical allodynia in the human masseter muscle, **101** 221

Taiminen, T., see Hagelberg, N., **101** 149
Takeda, D., Nakatsuka, T., Papke, R., Gu, J.G., Modulation of inhibitory synaptic activity by a non-$\alpha 4\beta 2$, non-$\alpha 7$ subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord, **101** 13
Tinazzi, M., see Rosso, T., **101** 117
Tokunaga, A., see Obata, K., **101** 65
Tomkins, E., see Cassidy, E.M., **101** 283
Tonali, P., see de Tommaso, M., **101** 25
Trempe, T.M., see Wacnik, P.W., **101** 175

Valeriani, M., see de Tommaso, M., **101** 25
van der Wouden, J.C., see Merlijn, V.P.B.M., **101** 33
Vierck Jr, C.J., see Staud, R., **101** 167
Vlaeyen, J.W.S., see Roelofs, J., **101** 299
Vowles, K.E., Gross, R.T., Work-related beliefs about injury and physical capability for work in individuals with chronic pain, **101** 291

Wacnik, P.W., Kehl, L.J., Trempe, T.M., Ramnaraine, M.L., Beitz, A.J., Wilcox, G.L., Tumor implantation in mouse humerus evokes movement-related hyperalgesia exceeding that evoked by intramuscular carrageenan, **101** 175
Wang, K., see Svensson, P., **101** 221
Warncke, T., see Jørum, E., **101** 229
Wei, H., see Pertovaara, A., **101** 237
Weintraub, W.S., see Kimble, L.P., **101** 45
Wilcox, G.L., see Wacnik, P.W., **101** 175
Wright, C.C., see Chesterton, L.S., **101** 259

Yamanaka, H., see Obata, K., **101** 65
Yi, D., see Obata, K., **101** 65
Yoshikawa, H., see Obata, K., **101** 65

Zandbergen, A.M.A.J., see Schasfoort, F.C., **101** 79
Zanette, G., see Rosso, T., **101** 117

PAIN

Pain 101 (2003) 311–314

www.elsevier.nl/locate/pain

Subject index Volume 101 (2003)

Activating transcription factor 3
Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats, **101** 65
Adenosine
Characterization of the analgesic actions of adenosine: comparison of adenosine and remifentanil infusions in patients undergoing major surgical procedures, **101** 129
Adolescents
Psychosocial factors associated with chronic pain in adolescents, **101** 33
Afferent inhibition
Analgesia from a peripherally active κ-opioid receptor agonist in patients with chronic pancreatitis, **101** 89
Aging
Age-related differences in endogenous pain modulation: a comparison of diffuse noxious inhibitory controls in healthy older and younger adults, **101** 155
Algometry
Gender differences in pressure pain threshold in healthy humans, **101** 259
Allodynia
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia, **101** 139
Ambulatory accelerometry
Impact of upper limb complex regional pain syndrome type 1 on everyday life measured with a novel upper limb-activity monitor, **101** 79
γ-Aminobutyric acidA-receptor
Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats, **101** 65
Analgesia
Reduction of TMD pain by high-frequency vibration: a spatial and temporal analysis, **101** 267
Antinociception
Powerful antinociceptive effects of the cone snail venom-derived subtype-selective NMDA receptor antagonists conantokins G and T, **101** 109
Vagus nerve stimulation in awake rats reduces formalin-induced nociceptive behaviour and fos-immunoreactivity in trigeminal nucleus caudalis, **101** 3
Brain plasticity
Functional plasticity in the human primary somatosensory cortex following acute lesion of the anterior lateral spinal cord: neurophysiological evidence of short-term cross-modal plasticity, **101** 117
Brain-derived neurotrophic factor
Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats, **101** 65
Breakthrough pain
Cancer breakthrough pain characteristics and responses to treatment at a VA medical center, **101** 55
Burning mouth syndrome
Striatal dopamine D1 and D2 receptors in burning mouth syndrome, **101** 149
Cancer
Immediate- or sustained-release morphine for dose finding during start of morphine to cancer patients: a randomized, double-blind trial, **101** 193
Cancer pain
Tumor implantation in mouse humerus evokes movement-related hyperalgesia exceeding that evoked by intramuscular carrageenan, **101** 175
Central pain
Thalamic field potentials in chronic central pain treated by periventricular gray stimulation – a series of eight cases, **101** 97
Central sensitisation
Reflexes in sympathetic vasoconstrictor neurones arising from urinary bladder afferents are not amplified early after inflammation in the anaesthetised cat, **101** 251
Powerful antinociceptive effects of the cone snail venom-derived subtype-selective NMDA receptor antagonists conantokins G and T, **101** 109
c-Fos
Vagus nerve stimulation in awake rats reduces formalin-induced nociceptive behaviour and fos-immunoreactivity in trigeminal nucleus caudalis, **101** 3
Chronic constriction injury
Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats, **101** 65
Chronic daily headache
Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
Chronic migraine
Abnormal brain processing of cutaneous pain in patients with chronic migraine, **101** 25
Chronic pain
Analgesia from a peripherally active κ-opioid receptor agonist in patients with chronic pancreatitis, **101** 89
Psychosocial factors associated with chronic pain in adolescents, **101** 33
Striatal dopamine D1 and D2 receptors in burning mouth syndrome, **101** 149
Work-related beliefs about injury and physical capability for work in individuals with chronic pain, **101** 291
Clinical trial
Analgesia from a peripherally active κ-opioid receptor agonist in patients with chronic pancreatitis, **101** 89
Cold allodynia/hyperalgesia
Cold allodynia and hyperalgesia in neuropathic pain: the effect of *N*-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo, **101** 229
Complex regional pain syndrome 1
Impact of upper limb complex regional pain syndrome type 1 on everyday life measured with a novel upper limb-activity monitor, **101** 79
Conantokins
Powerful antinociceptive effects of the cone snail venom-derived

doi:10.1016/S0304-3959(03)00035-6

subtype-selective NMDA receptor antagonists conantokins G and T, **101** 109
Cone snail venom-derived peptides
Powerful antinociceptive effects of the cone snail venom-derived subtype-selective NMDA receptor antagonists conantokins G and T, **101** 109
Congenital indifference
Congenital insensitivity to pain: an update, **101** 213
Congenital insensitivity
Congenital insensitivity to pain: an update, **101** 213
Cordotomy
Functional plasticity in the human primary somatosensory cortex following acute lesion of the anterior lateral spinal cord: neurophysiological evidence of short-term cross-modal plasticity, **101** 117
Cross-modal plasticity
Functional plasticity in the human primary somatosensory cortex following acute lesion of the anterior lateral spinal cord: neurophysiological evidence of short-term cross-modal plasticity, **101** 117
Deep brain stimulation
Thalamic field potentials in chronic central pain treated by periventricular gray stimulation – a series of eight cases, **101** 97
Delta opioid receptor
Up-regulation and trafficking of δ opioid receptor in a model of chronic inflammation: implications for pain control, **101** 199
Descending pain modulation
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Diffuse noxious inhibitory control
Age-related differences in endogenous pain modulation: a comparison of diffuse noxious inhibitory controls in healthy older and younger adults, **101** 155
Diffuse noxious inhibitory controls
Diffuse noxious inhibitory controls (DNIC) attenuate temporal summation of second pain in normal males but not in normal females or fibromyalgia patients, **101** 167
Disability
Work-related beliefs about injury and physical capability for work in individuals with chronic pain, **101** 291
Dopamine receptor
Striatal dopamine D1 and D2 receptors in burning mouth syndrome, **101** 149
Dorsal root ganglion
Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats, **101** 65
Double-blind method
Effects on muscle pain by intramuscular injection of granisetron in patients with fibromyalgia, **101** 275
Elderly
Age-related differences in endogenous pain modulation: a comparison of diffuse noxious inhibitory controls in healthy older and younger adults, **101** 155
Electron microscopy
Up-regulation and trafficking of δ opioid receptor in a model of chronic inflammation: implications for pain control, **101** 199
Experimental pain
Gender differences in pressure pain threshold in healthy humans, **101** 259
Factor analysis
The pain vigilance and awareness questionnaire (PVAQ): further psychometric evaluation in fibromyalgia and other chronic pain syndromes, **101** 299
Fear-avoidance
Work-related beliefs about injury and physical capability for work in individuals with chronic pain, **101** 291
Fibromyalgia
Diffuse noxious inhibitory controls (DNIC) attenuate temporal summation of second pain in normal males but not in normal females or fibromyalgia patients, **101** 167
Effects on muscle pain by intramuscular injection of granisetron in patients with fibromyalgia, **101** 275
Fibrosarcoma
Tumor implantation in mouse humerus evokes movement-related hyperalgesia exceeding that evoked by intramuscular carrageenan, **101** 175
Forelimb
Tumor implantation in mouse humerus evokes movement-related hyperalgesia exceeding that evoked by intramuscular carrageenan, **101** 175
Formalin test
Vagus nerve stimulation in awake rats reduces formalin-induced nociceptive behaviour and fos-immunoreactivity in trigeminal nucleus caudalis, **101** 3
Gamma-aminobutyric acid
Modulation of inhibitory synaptic activity by a non-α4β2, non-α7 subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord, **101** 13
Gender
Glutamate-evoked pain and mechanical allodynia in the human masseter muscle, **101** 221
Gender differences
Gender differences in pain characteristics of chronic stable angina and perceived physical limitation in patients with coronary artery disease, **101** 45
Gender differences in pressure pain threshold in healthy humans, **101** 259
Glycine
Modulation of inhibitory synaptic activity by a non-α4β2, non-α7 subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord, **101** 13
Hereditary sensory and autonomic neuropathy
Congenital insensitivity to pain: an update, **101** 213
5-HT
Effects on muscle pain by intramuscular injection of granisetron in patients with fibromyalgia, **101** 275
5HT1A
Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
5-HT-antagonist
Effects on muscle pain by intramuscular injection of granisetron in patients with fibromyalgia, **101** 275
Hyperalgesia
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Reflexes in sympathetic vasoconstrictor neurones arising from urinary bladder afferents are not amplified early after inflammation in the anaesthetised cat, **101** 251
Immediate-release morphine
Immediate- or sustained-release morphine for dose finding during start of morphine to cancer patients: a randomized, double-blind trial, **101** 193
Inflammation
Up-regulation and trafficking of δ opioid receptor in a model of chronic inflammation: implications for pain control, **101** 199
Inhibitory postsynaptic currents
Modulation of inhibitory synaptic activity by a non-α4β2, non-α7 subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord, **101** 13
Intrathecal catheterization
Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia, **101** 139
Laser-evoked potentials
Abnormal brain processing of cutaneous pain in patients with chronic migraine, **101** 25

Lower limb somatosensory evoked potentials
Functional plasticity in the human primary somatosensory cortex following acute lesion of the anterior lateral spinal cord: neurophysiological evidence of short-term cross-modal plasticity, **101** 117
Masseter muscle
Glutamate-evoked pain and mechanical allodynia in the human masseter muscle, **101** 221
McGill pain questionnaire – short form
Gender differences in pain characteristics of chronic stable angina and perceived physical limitation in patients with coronary artery disease, **101** 45
Mechanical allodynia/hyperalgesia
Cold allodynia and hyperalgesia in neuropathic pain: the effect of *N*-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo, **101** 229
Microdialysis
Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia, **101** 139
Migraine
Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
Mobility
Impact of upper limb complex regional pain syndrome type 1 on everyday life measured with a novel upper limb-activity monitor, **101** 79
Modeling
Psychosocial factors associated with chronic pain in adolescents, **101** 33
Morphine
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Immediate- or sustained-release morphine for dose finding during start of morphine to cancer patients: a randomized, double-blind trial, **101** 193
Tumor implantation in mouse humerus evokes movement-related hyperalgesia exceeding that evoked by intramuscular carrageenan, **101** 175
Movement-related hyperalgesia
Tumor implantation in mouse humerus evokes movement-related hyperalgesia exceeding that evoked by intramuscular carrageenan, **101** 175
Musculoskeletal pain
Tumor implantation in mouse humerus evokes movement-related hyperalgesia exceeding that evoked by intramuscular carrageenan, **101** 175
Neuropathic pain
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Cold allodynia and hyperalgesia in neuropathic pain: the effect of *N*-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo, **101** 229
Pain phenomena and possible mechanisms in patients with painful polyneuropathy, **101** 187
Powerful antinociceptive effects of the cone snail venom-derived subtype-selective NMDA receptor antagonists conantokins G and T, **101** 109
Neuropathy
Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia, **101** 139
Neuroticism
Psychosocial factors associated with chronic pain in adolescents, **101** 33
Nicotinic receptors
Modulation of inhibitory synaptic activity by a non-α4β2, non-α7 subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord, **101** 13
N-methyl-D-aspartate receptor
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
N-methyl-D-aspartate-receptor antagonist ketamine
Cold allodynia and hyperalgesia in neuropathic pain: the effect of *N*-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo, **101** 229
Noxious
Age-related differences in endogenous pain modulation: a comparison of diffuse noxious inhibitory controls in healthy older and younger adults, **101** 155
Ongoing pain
Cold allodynia and hyperalgesia in neuropathic pain: the effect of *N*-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo, **101** 229
Opioid agonist alfentanil
Cold allodynia and hyperalgesia in neuropathic pain: the effect of *N*-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo, **101** 229
Pain interference
Cancer breakthrough pain characteristics and responses to treatment at a VA medical center, **101** 55
Pain modulation
Age-related differences in endogenous pain modulation: a comparison of diffuse noxious inhibitory controls in healthy older and younger adults, **101** 155
Pain syndromes
Cancer breakthrough pain characteristics and responses to treatment at a VA medical center, **101** 55
Pain threshold
Effects on muscle pain by intramuscular injection of granisetron in patients with fibromyalgia, **101** 275
Pain vigilance and awareness questionnaire
The pain vigilance and awareness questionnaire (PVAQ): further psychometric evaluation in fibromyalgia and other chronic pain syndromes, **101** 299
Pain
Age-related differences in endogenous pain modulation: a comparison of diffuse noxious inhibitory controls in healthy older and younger adults, **101** 155
Analgesia from a peripherally active κ-opioid receptor agonist in patients with chronic pancreatitis, **101** 89
Congenital insensitivity to pain: an update, **101** 213
Functional plasticity in the human primary somatosensory cortex following acute lesion of the anterior lateral spinal cord: neurophysiological evidence of short-term cross-modal plasticity, **101** 117
Immediate- or sustained-release morphine for dose finding during start of morphine to cancer patients: a randomized, double-blind trial, **101** 193
Up-regulation and trafficking of δ opioid receptor in a model of chronic inflammation: implications for pain control, **101** 199
Pain-related behavior
Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats, **101** 65
Parents
Psychosocial factors associated with chronic pain in adolescents, **101** 33
Patch-clamp technique
Modulation of inhibitory synaptic activity by a non-α4β2, non-α7 subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord, **101** 13
Peers
Psychosocial factors associated with chronic pain in adolescents, **101** 33
Periaqueductal gray
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Periventricular gray
Thalamic field potentials in chronic central pain treated by periventricular gray stimulation – a series of eight cases, **101** 97
PET
Striatal dopamine D1 and D2 receptors in burning mouth syndrome, **101** 149

Physical functional limitation
Gender differences in pain characteristics of chronic stable angina and perceived physical limitation in patients with coronary artery disease, **101** 45
Postoperative pain
Characterization of the analgesic actions of adenosine: comparison of adenosine and remifentanil infusions in patients undergoing major surgical procedures, **101** 129
Pressure pain threshold
Gender differences in pressure pain threshold in healthy humans, **101** 259
Prolactin
Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
Prostaglandins
Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia, **101** 139
Quality of life
Gender differences in pain characteristics of chronic stable angina and perceived physical limitation in patients with coronary artery disease, **101** 45
Receptor trafficking
Up-regulation and trafficking of δ opioid receptor in a model of chronic inflammation: implications for pain control, **101** 199
Reinforcement
Psychosocial factors associated with chronic pain in adolescents, **101** 33
Reliability
The pain vigilance and awareness questionnaire (PVAQ): further psychometric evaluation in fibromyalgia and other chronic pain syndromes, **101** 299
Remifentanil
Characterization of the analgesic actions of adenosine: comparison of adenosine and remifentanil infusions in patients undergoing major surgical procedures, **101** 129
Rostroventromedial medulla
A dissociative change in the efficacy of supraspinal versus spinal morphine in the neuropathic rat, **101** 237
Sensory interaction
Reduction of TMD pain by high-frequency vibration: a spatial and temporal analysis, **101** 267
Sensory testing
Pain phenomena and possible mechanisms in patients with painful polyneuropathy, **101** 187
Serotonin
Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
Signs
Pain phenomena and possible mechanisms in patients with painful polyneuropathy, **101** 187
Somatosensory cortex
Functional plasticity in the human primary somatosensory cortex following acute lesion of the anterior lateral spinal cord: neurophysiological evidence of short-term cross-modal plasticity, **101** 117
Spinal cord
Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia, **101** 139
Spinal cord slice preparation
Modulation of inhibitory synaptic activity by a non-α4β2, non-α7 subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord, **101** 13
Start
Immediate- or sustained-release morphine for dose finding during start of morphine to cancer patients: a randomized, double-blind trial, **101** 193
Stress
Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
Sustained-release morphine
Immediate- or sustained-release morphine for dose finding during start of morphine to cancer patients: a randomized, double-blind trial, **101** 193
Sympathetic reflexes
Reflexes in sympathetic vasoconstrictor neurones arising from urinary bladder afferents are not amplified early after inflammation in the anaesthetised cat, **101** 251
Symptoms
Pain phenomena and possible mechanisms in patients with painful polyneuropathy, **101** 187
Temporomandibular disorders
Glutamate-evoked pain and mechanical allodynia in the human masseter muscle, **101** 221
Reduction of TMD pain by high-frequency vibration: a spatial and temporal analysis, **101** 267
Thalamic field potentials
Thalamic field potentials in chronic central pain treated by periventricular gray stimulation – a series of eight cases, **101** 97
Transformed
Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone, **101** 283
Trigeminal nucleus caudalis
Vagus nerve stimulation in awake rats reduces formalin-induced nociceptive behaviour and fos-immunoreactivity in trigeminal nucleus caudalis, **101** 3
Trigeminal pain
Glutamate-evoked pain and mechanical allodynia in the human masseter muscle, **101** 221
Upper limb activity
Impact of upper limb complex regional pain syndrome type 1 on everyday life measured with a novel upper limb-activity monitor, **101** 79
Urinary bladder
Reflexes in sympathetic vasoconstrictor neurones arising from urinary bladder afferents are not amplified early after inflammation in the anaesthetised cat, **101** 251
Vagus nerve stimulation
Vagus nerve stimulation in awake rats reduces formalin-induced nociceptive behaviour and fos-immunoreactivity in trigeminal nucleus caudalis, **101** 3
Validity
The pain vigilance and awareness questionnaire (PVAQ): further psychometric evaluation in fibromyalgia and other chronic pain syndromes, **101** 299
Vibration
Reduction of TMD pain by high-frequency vibration: a spatial and temporal analysis, **101** 267
Visceral afferents
Reflexes in sympathetic vasoconstrictor neurones arising from urinary bladder afferents are not amplified early after inflammation in the anaesthetised cat, **101** 251
Visceral pain
Analgesia from a peripherally active κ-opioid receptor agonist in patients with chronic pancreatitis, **101** 89
Reflexes in sympathetic vasoconstrictor neurones arising from urinary bladder afferents are not amplified early after inflammation in the anaesthetised cat, **101** 251
Visual analog pain scale
Effects on muscle pain by intramuscular injection of granisetron in patients with fibromyalgia, **101** 275
Wind-up
Diffuse noxious inhibitory controls (DNIC) attenuate temporal summation of second pain in normal males but not in normal females or fibromyalgia patients, **101** 167
Work capability
Work-related beliefs about injury and physical capability for work in individuals with chronic pain, **101** 291